

## Product datasheet

# Anti-ARID1A antibody [EPR13501] ab182560

**KO VALIDATED** Recombinant RabMAb

[11 References](#) [5 Images](#)

### Overview

<b>Product name</b>	Anti-ARID1A antibody [EPR13501]
<b>Description</b>	Rabbit monoclonal [EPR13501] to ARID1A
<b>Host species</b>	Rabbit
<b>Tested applications</b>	<b>Suitable for:</b> Flow Cyt, IHC-P, ICC/IF
<b>Species reactivity</b>	<b>Reacts with:</b> Mouse, Rat, Human
<b>Immunogen</b>	Recombinant fragment within Human ARID1A aa 1200-1350. The exact sequence is proprietary. Database link: <a href="#">O14497</a>
<b>Positive control</b>	IHC-P: Human kidney and human adenocarcinoma of endometrium without ARID1A mutation tissues. ICC/IF: Wildtype HAP1 and SH-SY5Y cells.
<b>General notes</b>	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> <li>- High batch-to-batch consistency and reproducibility</li> <li>- Improved sensitivity and specificity</li> <li>- Long-term security of supply</li> <li>- Animal-free production</li> </ul> <p>For more information <a href="#">see here</a>.</p> <p>Our RabMAb<sup>®</sup> technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to <a href="#">RabMAb<sup>®</sup> patents</a>.</p>

### Properties

<b>Form</b>	Liquid
<b>Storage instructions</b>	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
<b>Storage buffer</b>	pH: 7.20 Preservative: 0.01% Sodium azide Constituents: 40% Glycerol, 59% PBS, 0.05% BSA
<b>Purity</b>	Protein A purified
<b>Clonality</b>	Monoclonal
<b>Clone number</b>	EPR13501

Isotype

IgG

## Applications

Our [Abpromise guarantee](#) covers the use of **ab182560** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
Flow Cyt		Use at an assay dependent concentration. <a href="#">ab172730</a> - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.
IHC-P		1/1000. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.
ICC/IF		1/500.

## Target

### Function

Involved in transcriptional activation and repression of select genes by chromatin remodeling (alteration of DNA-nucleosome topology). Binds DNA non-specifically. Also involved in vitamin D-coupled transcription regulation via its association with the WINAC complex, a chromatin-remodeling complex recruited by vitamin D receptor (VDR), which is required for the ligand-bound VDR-mediated transrepression of the CYP27B1 gene. Belongs to the neural progenitors-specific chromatin remodeling complex (npBAF complex) and the neuron-specific chromatin remodeling complex (nBAF complex). During neural development a switch from a stem/progenitor to a post-mitotic chromatin remodeling mechanism occurs as neurons exit the cell cycle and become committed to their adult state. The transition from proliferating neural stem/progenitor cells to post-mitotic neurons requires a switch in subunit composition of the npBAF and nBAF complexes. As neural progenitors exit mitosis and differentiate into neurons, npBAF complexes which contain ACTL6A/BAF53A and PHF10/BAF45A, are exchanged for homologous alternative ACTL6B/BAF53B and DPF1/BAF45B or DPF3/BAF45C subunits in neuron-specific complexes (nBAF). The npBAF complex is essential for the self-renewal/proliferative capacity of the multipotent neural stem cells. The nBAF complex along with CREST plays a role regulating the activity of genes essential for dendrite growth.

### Tissue specificity

Highly expressed in spleen, thymus, prostate, testis, ovary, small intestine, colon, and PBL, and at a much lower level in heart, brain, placenta, lung, liver, skeletal muscle, kidney, and pancreas.

### Sequence similarities

Contains 1 ARID domain.


### Post-translational modifications

Phosphorylated upon DNA damage, probably by ATM or ATR.

### Cellular localization

Nucleus.


## Images



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-ARID1A antibody [EPR13501] (ab182560)


Immunohistochemical analysis of paraffin embedded Human kidney tissue (Left image) labeling ARID1A using ab182560 at 1/1000 dilution. Right image: Right picture: paraffine embedded human clear cell carcinoma of kidney with ARID1A mutation. A Ready to use HRP Polymer for Rabbit IgG (prediluted) was used as secondary. Counterstain: Hematoxylin.

Heat mediated antigen retrieval with Tris-EDTA, pH 9 was performed before commencing with IHC staining protocol.




Immunocytochemistry/ Immunofluorescence - Anti-ARID1A antibody [EPR13501] (ab182560)

ab182560 staining ARID1A in wild-type HAP1 cells (top panel) and ARID1A knockout HAP1 cells (bottom panel). The cells were fixed with 4% formaldehyde (10min), permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated with ab182560 at 1/500 dilution and ab195889 at 1/250 dilution (shown in pseudo colour red) overnight at +4°C, followed by a further incubation at room temperature for 1h with a goat secondary antibody to Rabbit IgG (Alexa Fluor® 488) (ab150081) at 2 µg/ml (shown in green). Nuclear DNA was labelled in blue with DAPI.




Flow Cytometry - Anti-ARID1A antibody [EPR13501] (ab182560)

Flow Cytometry analysis of SH-SY5Y (human neuroblastoma) cells labeling ARID1A with purified ab182560 at 1/230 dilution(10ug/ml) (red). Cells were fixed with 4% paraformaldehyde and permeabilised with 90% methanol. A Goat anti rabbit IgG (Alexa Fluor® 488)(1/2000 dilution) was used as the secondary antibody. Rabbit monoclonal IgG (Black) was used as the isotype control, cells without incubation with primary antibody and secondary antibody (Blue) were used as the unlabeled control.



Immunocytochemistry/ Immunofluorescence - Anti-ARID1A antibody [EPR13501] (ab182560)

Immunofluorescent analysis of SH-SY5Y cells labeling ARID1A with ab182560 at 1/500 and Goat anti rabbit IgG(Alexa Fluor®555) at 1/200. Image at the right stained with DAPI.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-ARID1A antibody [EPR13501] (ab182560)

Immunohistochemical analysis of paraffin embedded Human adenocarcinoma of endometrium without ARID1A mutation (Left image) labeling ARID1A using ab182560 at 1/1000 dilution. Right image: Right picture: paraffine embedded human adenocarcinoma of endometrium with ARID1A mutation. A Ready to use HRP Polymer for Rabbit IgG (prediluted) was used as secondary. Counterstain: Hematoxylin.

Heat mediated antigen retrieval with Tris-EDTA, pH 9 was performed before commencing with IHC staining protocol.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.com/abpromise> or contact our technical team.

### Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors